



Directed Donation Interval Less than 8 Weeks

Physician's permission for a person to serve as a blood donor more than once in 8 weeks¹

Donor Name _____

Date of Birth _____

I have examined the person listed above and found him/her to be in good health as indicated in part by:

1. Normal temperature
2. Systolic and diastolic blood pressure are within normal limits
3. Freedom from acute respiratory diseases
4. Freedom from any infectious skin disease at the site of phlebotomy and from any such disease generalized to such an extent as to create a risk of contamination of blood
5. Freedom from any disease transmissible by blood transfusion, insofar as can be determined by history and examinations indicated above, and
6. Freedom of arms and forearms from skin punctures or scars indicative of drug abuse.

The person may serve as a blood donor more than once in 8 weeks only if he/she meets other Blood Systems donor qualification criteria, including a blood hemoglobin level not less than 12.5 gm/100 mL of blood. I discussed the possible risk of donating frequently to include the possibility of iron store depletion.

Physician Name _____

Physician Signature _____

Date² _____

Address _____

Telephone _____

¹ 21 CFR 640.3(f)

² Must be on the date of donation prior to phlebotomy

Donor Informed Consent for Donation Interval Less than 8 Weeks

I, _____ understand the hazards of
(Donor Name)

donating whole blood at an interval of less than the normally required 8 weeks, as explained to me by

_____ and consent to donate.

Donor Signature

Date